

Cyperus limiticola, a New Name for a Madagascan *Cyperus* (Cyperaceae)

Isabel Larridon*, Marc F. M. Reynders, and Paul Goetghebeur

Ghent University, Department of Biology, Research Group Spermatophytes, K. L. Ledeganckstraat 35, 9000 Gent, Belgium. *Corresponding author: isabel.larridon@ugent.be

ABSTRACT. There is a need to propose a new name in *Cyperus* L. for *C. paucispiculatus* Chermезon because the name currently used is an illegitimate later homonym of *C. paucispiculatus* Böckeler. A new name, *C. limiticola* Larridon & Reynders, is provided here.

Key words: Cyperaceae, *Cyperus*.

In 1884, Böckeler described *Cyperus paucispiculatus* Böckeler [= *Pycrus divulsus* (Ridley) C. B. Clarke]. Chermезon (1919) used *C. paucispiculatus* Chermезon for a new species from Madagascar and placed it in its own section, *Cyperus* sect. *Paucispiculati* Chermезon. The latter species does not belong to the genus *Pycrus* P. Beauvois, and so is not related to *P. divulsus*, although both species show a reduction in the number of spikelets. *Cyperus paucispiculatus* Chermезon is a later homonym of *C. paucispiculatus* Böckeler and is therefore illegitimate (McNeill et al., 2006: Art. 53.1). A new name, *C. limiticola* Larridon & Reynders, is proposed. The epithet, which means “inhabiting the edge,” refers to the natural surroundings in which the plant was collected (i.e., the edge of the forest).

Cyperus limiticola Larridon & Reynders, nom. nov.

Replaced name: *Cyperus paucispiculatus* Chermезon, Bull. Soc. Bot. France 66: 342. 1919 [1920], non *Cyperus paucispiculatus* Böckeler, 1884. TYPE: Madagascar. [Antananarivo]: Distr. Manjakandriana, lisière de la forêt à l'Est d'Ambatolaona, 1400 m, 11 Nov. 1912, R. Viguier & J.-H. Humbert 1256 (holotype, P 00450832; isotypes, P 00450833, P 00581793). Figure 1.

Caespitose perennial herb with a slender rhizome; culms filiform, 15–35 cm × ca. 0.5 mm, smooth. Leaves few, slightly shorter than the culms, ca.

0.5 mm wide, setaceous, semicylindrical, smooth, leaf sheaths fuscous. Lower involucral bract smooth, erect, continuing in the direction of the culm, up to 3–6 cm, setaceous; second involucral bract glumiform; anthela composed of 1 to 3 radii; radii 0–4 cm, erect, with short prophylls and 1 or 2 spikelets; spikelets compressed, ovate-lanceolate, ca. 5 × 2.5 mm, with 5 to 7 flowers; rachilla winged. Glumes slightly imbricate, ovate-lanceolate, shortly mucronate, ca. 3 mm, subrigid, the sides yellow-brown, with 2 or 3 obscure lateral nerves, midrib narrow, smooth, green, 3-nerved; stamens 3; anthers linear, connectives shortly setiferous; style deeply trifid. Achenes immature.

Discussion. Chermезon (1919) stated that, with its multinervate glumes and its setiferous anthers, this isolated species only approaches the sections *Cyperus* sect. *Diffusi* (Kunth) C. B. Clarke and *Cyperus* sect. *Debilissimi* Chermезon, from which it is, however, sufficiently distinct, by its inflorescence and its leaves, to deserve its own section. Molecular research may offer an answer to its taxonomic relationships, although sadly this species is only known from its immature type specimens.

Acknowledgments. We thank M. Pignal (P) for the loan of the specimens examined for this study.

Literature Cited

- Böckeler, O. 1884. Neue Cyperaceen. Bot. Jahrb. Syst. 5: 497–498.
Chermезon, H. 1919 [1920]. *Cyperus* nouveaux de Madagascar. Bull. Soc. Bot. France 66: 338–353.
McNeil, J., F. R. Barrie, H. M. Burdet, V. Demoulin, D. L. Hawksworth, K. Marhold, D. H. Nicolson, J. Prado, P. C. Silva, J. E. Skog, J. H. Wiersema & N. J. Turland (editors). 2006. International Code of Botanical Nomenclature (Vienna Code). Regnum Veg. 146.




Figure 1. *Cyperus limiticola* Larridon & Reynders. —A. Habit. —B. Inflorescence. —C. Detail of spikelets. Drawn by Marc Reynders from the holotype *Viguiier & Humbert 1256* (P).